

{Exhibit 35}

BEST AVAILABLE COPY

Langer and Ward, Abstract 1153: "A Rapid and Sensitive Immunological Method for In Situ Gene Mapping in Journal of Supramolecular Structure and Cellular Biology, (1981)

FILE COPY *Langer*

Journal of Supramolecular Structure and Cellular Biochemistry

SUPPLEMENT 5, 1981

COOPER, DUNCAN
GRIFPA & A.
RECEIVED

MAR 31 1981

J. P. WHITE

10th Annual
ICN-UCLA SYMPOSIUM
Molecular and Cellular Biology

Abstracts

FEBRUARY 8 – MARCH 20, 1981

BEST AVAILABLE COPY

Alan R. Liss, Inc., New York

Journal of Supramolecular Structure and Cellular Biochemistry

Editor:

C. FRED FOX, Molecular Biology Institute, University of California, Los Angeles, CA 90024

Associate Editors:

GIOVANNA FERRO-LUZZI AMES, Department of Biochemistry, University of California, Berkeley, Berkeley, CA 94720

RICHARD ANDERSON, Department of Cell Biology, University of Texas, Health Science Center, Dallas, TX 75230

J. KENT BLASIE, Department of Chemistry, University of Pennsylvania, Philadelphia, PA 19174

ARTHUR BLUME, Physiological Chemistry, Roche Institute of Molecular Biology, Nutley, NJ 07110

RALPH BRADSHAW, Biological Chemistry, Washington University, Medical School, St. Louis, MO 63110

DANIEL BRANTON, The Biological Laboratories, Harvard University, Cambridge, MA 02138

DENNIS D. CUNNINGHAM, Medical Microbiology, University of California, Irvine, Irvine, CA 92717

MICHAEL EDIDIN, Department of Biology, Johns Hopkins University, Baltimore, MD 21218

FREDERICK A. EISERLING, Department of Bacteriology, University of California, Los Angeles, CA 90024

ELLIOT ELSON, Biological Chemistry, Washington University, Medical School, St. Louis, MO 63110

WOLFGANG EPSTEIN, Department of Biochemistry, University of Chicago, Chicago, IL 60637

IAN GIBBONS, Pacific Biomedical Research Center, University of Hawaii, Honolulu, HI 96822

LUIS GLASER, Department of Biological Chemistry, Washington University School of Medicine, St. Louis, MO 63110

EUGENE GOLDWASSER, Department of Biochemistry, University of Chicago, Chicago, IL 60637

ULF HENNING, Max-Planck-Institut für Biologie, Corrensstrasse 38, 74 Tübingen, Germany

EDWARD KELLENBERGER, Biozentrum der Universität Basel, Klingelbergstrasse 70, CH-4056 Basel, Switzerland

JOHN H. LAW, Department of Biochemistry, University of Chicago, Chicago, IL 60637

HARVEY F. LODISH, Department of Biology, Massachusetts Institute of Technology, Cambridge, MA 02139

HARDEN M. MCCONNELL, Department of Chemistry, Stanford University, Stanford, CA 94305

GARTH L. NICOLSON, Department of Developmental and Cell Biology, University of California, Irvine, Irvine, CA 92717

DALE L. OXENDER, Department of Biological Chemistry, University of Michigan, Ann Arbor, MI 48109

ROBERT POYTON, Department of Microbiology, University of Connecticut, Farmington, CT 66032

JEAN PAUL REVEL, Division of Biology 156-29, California Institute of Technology, Pasadena, CA 91125

PHILLIPS W. ROBBINS, Massachusetts Institute of Technology, Cambridge, MA 02139

RUSSELL ROSS, Department of Pathology, University of Washington, Seattle, WA 98195

DAVID SABATINI, Cell Biology, New York University Medical School, New York, NY 10016

MILTON SAIER, Department of Biology, University of California, San Diego, La Jolla, CA 92037

ERIC SHOOTER, Department of Genetics, Stanford University, School of Medicine, Stanford, CA 94305

PHILIP SIEKEVITZ, The Rockefeller University, New York, NY 10021

ROBERT D. SIMONI, Department of Biology, Stanford University, Stanford, CA 94305

THEODORE L. STECK, Department of Biochemistry, University of Chicago, Chicago, IL 60637

EDWIN W. TAYLOR, Department of Biophysics and Theoretical Biology, University of Chicago, Chicago, IL 60637

GEORGE TODARO, Viral Leukemia & Lymphoma Branch, National Cancer Institute, NIH, Bethesda, MD 20205

JAMES WATSON, Department of Microbiology, University of California Irvine, Irvine, CA 92717

IRVING WEISSMAN, Department of Pathology, Stanford University, School of Medicine, Stanford, CA 94305

DAVID WILSON, Department of Biochemistry, Cornell University, Ithaca, NY 14853

H.G. WITTMANN, Max-Planck-Institut für Molekulare Genetik, Ihnestrasse 63-73, D-1000 Berlin 33 (Dahlem), Germany

Managing Editor, 1981 Abstract Supplement:

FRAN STUSSER, Molecular Biology Institute, University of California, Los Angeles, CA 90024

Journal of Supramolecular Structure and Cellular Biochemistry (ISSN 0091-7419) is published 13 times a year by Alan R. Liss, Inc., 150 Fifth Avenue, New York, NY 10011.

Subscription Price: Volumes 15, 16, and 17, plus Supplement 5, 1981 (thirteen issues): \$237.00 in US; \$271.00 in Europe, the Middle East, and Africa; \$260.00 in other countries. (Subscriptions going to Europe, the Middle East, and Africa will be sent airfreight to Europe and mailed from there.) Payments must be made in US dollars or at the current rate of exchange. For payments not made in US dollars and through a US bank, add \$6 service charge. Change of Address: Please send to publisher six weeks prior to move; enclose present mailing label with change of address. Claims for Missing Issues: Claims cannot be honored beyond four months after mailing date. Duplicate copies cannot be sent to replace issues not delivered because of failure to notify publisher of change of address. Exclusive agent in Japan: Igaku Shoin Limited, Foreign Department, 1-28-36 Hongo, Bunkyo-ku, Tokyo 113, Japan.

Printed in the United States of America.

Copyright © 1981 Alan R. Liss, Inc.

Supplement 5, 1981

Listed in Current Contents.

BEST AVAILABLE COPY

TUMOF
Edward
Arnold L

RATION
Bruce C

B and T
Ellen Vi

GENE F
Bert O'M

EVOLU
Ralph E
Gordon

INTERF
Thomas

1153

A RAPID AND SENSITIVE IMMUNOLOGICAL METHOD FOR IN SITU GENE MAPPING. Pennina R. Langer and David C. Ward, Department of Human Genetics, Yale University, New Haven, CT 06510. A method for in situ localization of specific DNA sequences has been developed which exploits the interaction between modified nucleotides and antibodies directed against the modification. Cloned, unique sequences of drosophila DNA are nick translated in vitro in the presence of E. Coli DNA Polymerase I and an analogue of dUTP which contains a biotin molecule covalently linked to the C5 position of the pyrimidine ring. The nick translated probe containing approximately 0.25% biotin substituted nucleotides, is hybridized in situ, according to standard protocols, to drosophila salivary gland chromosomes. After hybridization, the slides are incubated with monospecific Rabbit anti-biotin followed by FITC-Coat anti-rabbit IgG. After counterstaining with Evan's Blue, a single yellow-green fluorescent band, corresponding to the map location of the cloned DNA is seen against a red fluorescent background of the salivary gland chromosomes. Current work is aimed at further refining the system for use in localization of unique sequences on mammalian metaphase chromosomes. FITC-Avidin and histochemical reagents are under investigation as substitutes for the indirect immunofluorescent localization of hybridized regions. Other potential applications of biotinyl polynucleotides as affinity reagents will also be presented.

1154 DEVELOPMENTAL REGULATION OF SEA URCHIN HISTONE GENES IN XENOPUS EGGS AND OOCYTES. Laurence Etkin, Dept. of Zoology, University of Tennessee, Knoxville, TN 37916. Sea urchin histone genes H1, H4, and H2B, contained in the recombinant plasmid pSp102, were microinjected into either the germinal vesicle of oocytes or the cytoplasm of fertilized eggs of Xenopus laevis. Recipient oocytes were treated with progesterone, to induce maturation, and incubated in ^3H lysine to label newly synthesized histones. Histone proteins were extracted with 10% GuCl-40% EtOH and analyzed by two dimensional gel electrophoresis and fluorography. Newly synthesized sea urchin H1, and H2B histones were detected in recipient oocytes cultured without progesterone. These proteins are not observed, however, in recipient oocytes which have matured in response to progesterone. This suggests that the sea urchin histone genes are not expressed in oocytes undergoing maturation. When the plasmid pSp102 is injected into fertilized eggs shortly after artificial insemination, sea urchin histones are not detected until the late gastrula-early neurula stage of development, and continue to be expressed as late as the tailbud stage. These results indicate that the microinjected sea urchin histone genes are expressed concurrently with endogenous Xenopus histone genes in growing oocytes, matured oocytes, and embryos, suggesting that the sea urchin genes are able to recognize and respond to endogenous Xenopus gene regulatory signals.

1155 USE OF RECOMBINANT DNA MOLECULES TO INVESTIGATE REGULATORY SIGNALS FOR THE INITIATION OF TRANSCRIPTION, Peter Gruss and George Khoury, National Institutes of Health, Bethesda, MD 20205. Nucleotide signals required for the initiation of transcription for the early region of the SV40 genome have been identified. This has been achieved by deleting portions of the genome between 0.69 and 0.71 map unit. A 72 bp repeat has been found to harbor a cis essential function, which cannot be provided in trans by either the early tsA28 or the late tsB4 mutant. Further studies have shown that this mutant is incapable of inducing early viral RNA as examined by S1 nuclease analysis, or the early gene product, T-antigen. Studies are in progress to determine whether this mutant can replicate its DNA in the presence of exogenously provided T-antigen. Further attempts to characterize this repetitive element involve replacing the SV40 72 bp repeats by a related repeat unit derived from Moloney sarcoma virus. Results of these experiments will be discussed in detail.